



PDPM	PDPM				
Nursing	Function	Descriptors for			
Component	Score	Extensive Services	Requirements	ICD-10	MDS 3.0 Section
	If patients function score is 15 or 16,	Tracheostomy Care, Ventilator Respirator and Isolation for active infectious disease while	Nursing Function Score less than or equal to 14		O0100E2
	classify to Clinically Complex	a resident			O0100F2
					O0100M2
ES3	0-14		<ul> <li>Tracheostomy care (while a resident)</li> </ul> AND		O0100E2
			Ventilator/Respirator (while a resident)		O0100F2
ES2	0-14		<ul> <li>Tracheostomy care (while a resident)</li> <li>OR</li> </ul>		O0100E2
			Ventilator/Respirator (while a resident)		O0100E2
ES1	0-14		<ul> <li>Isolation for active infectious disease</li> <li>Without tracheostomy care</li> <li>And Without Ventilator/ Respirator care (while a patient)</li> </ul>		O0100M2





Nursing Function Descriptors for	PDPM	PDPM				
Component   Score   Special Care High   Requirements   ICD-10   MDS 3.0 Section			Descriptors for			
If patients nursing function score is 15 or 16, classify to Clinically Complex  HDE2 HDE1  O-5 O-5 O-6 HBC2 HBC1  HBC2 HBC1  HBC2 HBC1  HBC2 HBC1  HBC2 HBC1  Respiratory Therapy  If patients nursing functions core less than or equal to 14  Comatose, Septicemia, Respiratory Therapy  Espticemia, Respiratory Therapy  Ocaratose and completely ADL dependent  Septicemia  Diabetes with both daily injections (all 7 days) requiring physician order changes on 2 or more days (N0350B)  Quadriplegia and Nursing Function Score <=11  Chronic obstructive pulmonary disease and shortness of breath when lying flat  Fever with O Pneumonia O Vorniting O Feeding tube: 26-50% or more calories and at least 501 cc of fluid or 51% of the calories for the entire 7 days O Weight loss (K0300)  Parenteral/IV feedings Respiratory therapy for 7 days	_			Requirements	ICD-10	MDS 3.0 Section
ADL less than or equal to 14  Notes: Signs of depression used for end splits; PHQ-9 Score =>10  D0300 grerater than or equal to 10 but not 99  Or  D0600 greater than or equal	HDE2 HDE1	Score  If patients nursing function score is 15 or 16, classify to Clinically Complex  0-5 0-5	Special Care High Serious Medical Conditions: Comatose, Septicemia,	Nursing Function Score less than or equal to 14  Comatose and completely ADL dependent Septicemia Diabetes with both daily injections (all 7 days) requiring physician order changes on 2 or more days (N0350B) Quadriplegia and Nursing Function Score <=11 Chronic obstructive pulmonary disease and shortness of breath when lying flat Fever with Pneumonia Vomiting Feeding tube: 26-50% or more calories and at least 501 cc of fluid or 51% of the calories for the entire 7 days Weight loss (K0300) Parenteral/IV feedings Respiratory therapy for 7 days  AND ADL less than or equal to 14  Notes: Signs of depression used for end splits; PHQ-9 Score =>10  D0300 grerater than or equal to 10 but not 99	ICD-10	I, I2100  I, I2900; Section N, N0350,A  I, I5100  I, I6200  J, J1550,A I, I2000 J, J1550,B K, K0500,B K, K0300  K, K0500,A





PDPM Nursing Component	PDPM Function Score	Descriptors for Special Care Low	Requirements	ICD-10	MDS 3.0 Section
LDE2 LDE1	0-5 0-5	Serious Medical Conditions: Radiation Therapy or Dialysis	<ul> <li>Cerebral Palsy and Nursing         Function Score &lt;=11</li> <li>Multiple Sclerosis and Nursing         Function Score &lt;=11</li> <li>Parkinson's Disease and         Nursing Function Score &lt;=11</li> <li>Feeding tube 26-50% or more         calories and at least 501 cc of</li> </ul>		I, I4400 I, I5200 I, I5300 K, K0700,A,B,
LBC2 LBC1	6-14 6-14		fluid or 51% of the calories for the entire 7 days  Ulcers with 2 or more skin treatments  2 or more stage II  1 or more stage III or IV pressure ulcers  Unstageable secondary to slough/ eschar  2 or more venous/arterial ulcers OR  1 stage II pressure ulcer AND 1 venous/arterial ulcer  Foot infection, diabetic foot		M, M0800,A M, M0800,B,C M, M1030 M, M-0800,A M, M1030 M, M1040,A,C  O, 00100,B,2 I, I6300 O,00100C O, 00100,J,2
			ulcer or other open lesions on the foot with treatment  Radiation therapy while a resident  Dialysis while a resident  Notes: Signs of depression used for end splits; PHQ-9 Score =>10  D0300 grerater than or equal to 10 but not 99  Or		
			D0600 greater than or equal to 10		





PDPM Nursing	PDPM Function	Descriptors for			
Component	Score	Clinically Complex	Requirements	ICD-10	MDS 3.0 Section
CDE2 CDE1	0-5 0-5	Conditions requiring Complex Medical Care:	<ul> <li>Residents with Extensive         Services, Special Care High or         Special Care Low qualifier.         AND</li> <li>ADL score = 0-1</li> </ul>		
CBC2	6-14	Pneumonia, Surgical Wounds, Burns	OR  Residents with any one of the following Clinically Complex qualifiers:		I, I2000 I, I4900 M, M1040,E
CBC1	6-14		<ul> <li>Pneumonia</li> <li>Hemiplegia/Hemiparesis with Nursing Function Score &lt;=11</li> </ul>		M, M1040,F O, O0100,A,2 O, O0100,C,2
CA2 CA1	15-16 15-16		<ul> <li>Open Lesions (other than ulcers, rashes, and cuts) with any selected skin treatment or surgical wounds</li> <li>Burns</li> <li>Chemotherapy while a</li> </ul>		O, O0100,H,2 O, O0100,I,2
			resident  Oxygen while a resident  IV medications while a resident  Transfusions while a resident		
			Notes: Signs of depression used for end splits; PHQ-9 Score =>10		
			D0300 grerater than or equal to 10 but not 99		
			Or		
			D0600 greater than or equal to 10		





PDPM	PDPM	Descriptors for			
Nursing	Function	Behavioral or Cognitive			
Component	Score	Symptoms	Requirements	ICD-10	MDS 3.0 Section
	If patients	, ,	Nursing Function Score is 11 or	ICD-10	IVIDS 5.0 Section
	function		greater		C1000
	score is	Behavioral or Cognitive	greater		B0700
	less than	Symptoms	Cognitive impairment		B0700
	11, go to	Symptoms	BIMS score <= 9 or		E0100A E0100B
	reduced		<ul><li>CPS &gt;= 3</li></ul>		LOTOGA LOTOGB
	Physical		<ul><li>BIMS 15, best score,</li></ul>		
	Function		cognitively intact		
'	runction		BIMS 0, worst		
			performance		
BAB2	11-16		BIMS 99, interview not		
5, 152			successful		
			3466633141		E, E0200,A,B,C
			OR		E, E0300,1
BAB1	11-16				, ,
			Hallucinations or delusions		E, E0900
			OR		•
			<b></b>		
			Residents displaying any of the		
			following on 4 or more days over		
			last 7 days:		
			·		
			Physical or verbal behavioral		
			symptoms toward others		
			Other behavioral symptoms		
			Rejection of care		
			Wandering		
			AND		
			ADL score <=5		
			ADL SCOTE <=3		
			Notes: Restorative Nursing used		
			for end splits. See Reduced		
			Physical Function for Restorative		
			Nursing Services count		
			5		





PDPM Nursing Component	PDPM Function Score	Descriptors for Reduced Physical Function	Requirements	ICD-10	MDS 3.0 Section
PDE2 PDE1  PBC2 PBC1  PA2 PA1	0-5 0-5 6-14 6-14 15-16 15-16	Assistance with Daily Living and General Supervision	Residents not qualifying for other categories  Restorative Nursing Services: Urinary and/or bowel training program Passive and/or active ROM Amputation/prosthesis care training Splint or brace assistance Dressing or grooming training Eating or swallowing training Transfer training Bed mobility and/or walking training Communication training		O, 0500,A-J H, H0200/H0500